

The Baltimore Traffic Study: Assessment of Air Quality Impact and Model Development

Timothy J. Buckley

Alison Geyh

Patrick N. Breysse

Johns Hopkins Bloomberg School of Public Health
Baltimore, MD


**National Training Workshop On Local Urban Air Toxics Assessment
And Reduction Strategies
November 13 – 15, 2001
Detroit, MI**


Purpose

- Assess the impact of traffic on community air quality
- Evaluate traffic related air quality variability by hour of day, day of week, season
- Characterize relationship between indoor and outdoor air pollution concentrations

The Community

- Midtown
 - 100 City Blocks
 - Population of 11,000
 - 25% with HH Income <\$15,000
 - Bolton Hill (36% AA)
 - Mount Vernon (35% AA)
 - Charles North (92% AA)
 - Transportation Impact
 - Four major arterials
 - Expressway
 - Rail line





Midtown Community Plan

Bolton Hill

Charles North

Madison Park

Mount Vernon

Chapter II. Changing the Environment

- “Midtown’s greatest problem is its current pattern of traffic, transit and parking . . .”
- “More than any other factor or combination of factors, existing conditions in transportation are weakening Midtown and preventing it from fulfilling its mission as a magnet for people and investment.”
- Traffic: There is too much traffic, particularly on key Residential streets like Calvert and St. Paul. Everyone of Midtown’s main north-south streets now serves as a commuter raceway and bus route.






Rationale

- Community / Public Health
 - Toxic emissions
 - Particles (Kleeman et al. ES&T)
 - Diesel (Kinney et al. 2000)
 - PAH (Dubowsky; Dickhut et al. 2000)
 - CO, NO_x, and VOCs (Chase et al. 2000)
 - Dioxins and furans (Ryan and Gullett, 1999)
 - Emissions in close proximity and integral to human populations
 - Health
 - Cancer (Pearson et al. 2000)
 - Respiratory effects in children (Brunekreef et al. 1997, Wjst et al. 1993, Weiland et al. 1994)

Impact of Traffic on Indoor PAH

Weekday




Geometric Mean (ng/m³)

Urban: 31*

Semi-Urban: 19*

Suburban: 8.0*

Weekend



Geometric Mean (ng/m³)

Urban: 10

Semi-Urban: 9.8

Suburban: 4.9*

Dubowsky SD; Wallace LA; Buckley TJ (1999): The contribution of traffic to indoor concentrations of polycyclic aromatic hydrocarbons. *J. Exposure Anal. Environ. Epi.* 9(4):312-320.

Outdoor

Indoor

41.5 lpm

(3-Story Row Home
located on the 1700 block
of North Calvert)

Support Instruments

TEI 146C Multi-
Gas Calibrator

TEI 111 Zero
Air Generator

ESC 8816
Data Logger

Manifold Diameter: 4.76 cm

Isokinetic Ports (@41.5 lpm)

60 cm

Outdoor Indoor
Aethalometer™

Dasibi 1003 AH
O₃ Monitor

TEI 42C NO_x
Monitor

Pump

EcoChem
2000 PAH
Climet
CI-500



TEI 55C
VOC Monitor

TEI 48C CO
Monitor

Sampling Manifold

Indicates unfunded

Instrumentation




Sampling Manifold

The Residential Fixed Site Monitoring Station



Weekday CO Concentration by Hour (Aug 2000 – Sep 2001)




(NAAQS: 1h=35 ppm; 8h=9 ppm)

*Draft, Do Not
Cite or Quote*

Indoor and Outdoor CO Concentration by Day of Week


*Draft, Do Not
Cite or Quote*



Draft, Do Not
Cite or Quote

Draft, Do Not
Cite or Quote


Weekday PAH Concentration by Hour



Draft, Do Not
Cite or Quote

PAH Measured Indoors and Outdoors by Day of Week


Draft, Do Not
Cite or Quote




Draft, Do Not
Cite or Quote

Weekday Black Carbon Concentration by Hour

Draft, Do Not
Cite or Quote




Black Carbon Measured Indoors and Outdoors by Day of Week



LaRosa et al. (in press): Rural median: $0.70 \mu\text{g}/\text{m}^3$; Urban Median: $1.4 \mu\text{g}/\text{m}^3$

Median Air Pollution by Day of Week and Location



Draft, Do Not
Cite or Quote

Correlation Coefficient for Total Data Set

Draft, Do Not
Cite or Quote

| | Indoor CO | Outdoor CO | Indoor PAH | Outdoor PAH | Indoor BC | Outdoor BC | Mean count |
|-------------|---|--|------------------------|------------------------|------------------------|------------------------|------------------|
| Indoor CO | 1.00 ^a b 9776 ^c | a : Pearson Correlation Coefficients b : Prob > r under H0 : Rho=0 c : Number of Observation | | | | | |
| Outdoor CO | 0.98 <.0001 9776 | 1.00 9776 | | | | | |
| Indoor PAH | 0.46 <.0001 9760 | 0.48 <.0001 9760 | 1.00 9760 | | | | |
| Outdoor PAH | 0.38 <.0001 9770 | 0.43 <.0001 9770 | 0.84 <.0001 9754 | 1.00 9770 | | | |
| Indoor BC | 0.28 <.0001 7426 | 0.30 <.0001 7426 | 0.72 <.0001 7425 | 0.59 <.0001 7423 | 1.00 7657 | | |
| Outdoor BC | 0.29 <.0001 7306 | 0.36 <.0001 7306 | 0.64 <.0001 7305 | 0.75 <.0001 7303 | 0.70 <.0001 7537 | 1.00 7537 | |
| Mean count | -0.08 <.0001 8566 | -0.03 0.00 8566 | 0.07 <.0001 8565 | 0.23 <.0001 8561 | 0.04 0.00 6799 | 0.18 <.0001 6679 | 1.00 9324 |

Conclusions

- Traffic-related community air quality impact evaluated for BC and PAH
- CO concentrations low and weak traffic effect
- BC and PAH Indoor/Outdoor ratios < 1 indicate penetration loss
- Future Work
 - Continued data analysis (seasonal variability, meteorology)
 - Biomarker study (benzene & 1-hydroxypyrene)
 - Community exposure study

Acknowledgements

- Sung Roul Kim for data analysis support
- Funding provided by the U.S. EPA